

KEK-TH-507
 YUMS-97-002
 SNUTP-97-009
 (February 1, 2008)

Is CDF large p_T anomaly from virtual SUSY threshold effects?*

C.S. Kim^{1,2}

1: *Department of Physics, Yonsei University, Seoul 120-749, Korea*

2: *Theory Group, KEK, Tsukuba, Ibaraki 305, Japan*

Abstract

Talk is given at the YITP International Workshop – ‘Recent Developments in QCD and Hadron Physics’. Recent CDF data of the inclusive jet cross section shows anomalous deviation around large transverse momentum $p_T(j) \approx 200 \sim 350$ GeV. Is it possible to interpret the anomaly in terms of virtual SUSY effects. The answer is ‘NO’, because we find that the virtual SUSY loop interference effects are too small to explain the CDF data.

I. INTRODUCTION

The CDF [1] and D0 [2] collaborations at Tevatron Collider have recently reported data for the inclusive jet cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1800$ GeV. Let us recapitulate some particulars of this data.

- (a) The CDF measurement is based on a data sample of 19.5 pb^{-1} collected in 1992-93 with the CDF detector at the Tevatron collider. Jets were reconstructed using a cone algorithm.

*Talk is given at the YITP International Workshop – ‘Recent Developments in QCD and Hadron Physics’, on Dec. 16–18, at Kyoto Japan. Proceedings will be published, edited by Y. Koike.

- (b) The measurements have been reported over a wide range of transverse energy, $15 \text{ GeV} \leq p_T \leq 440 \text{ GeV}$, and around the central pseudorapidity region $0.1 \leq |\eta| \leq 0.7$. The smallest distance probed is on the order of 10^{-19} m .
- (c) After accounting for uncertainties the corrected experimental cross section, when compared to the Next-to-leading [NLO] QCD predictions using MRSD0' parton distribution function [PDF's], is significantly higher than the NLO prediction for $p_T > 200 \text{ GeV}$. For p_T below 200 GeV the agreement between the CDF and the NLO QCD cross section is excellent, while the D0 results are higher than the NLO PQCD predictions within the statistical uncertainties.

There are basically two logical possibilities for explanations of the CDF data on inclusive jet production cross sections: (i) the parton distribution functions determined at low Q^2 region may not be accurate enough to be applicable to the high p_T region with $p_T > 200 \text{ GeV}$, or (ii) there are some new physics around the electroweak scale. We give a brief review of these two possibilities in the following.

The most interesting possibility is that jet measurements at hadron colliders may be sensitive to quantum corrections due to virtual SUSY particles [3,4,5]. The purpose of this note is to concentrate on this scenario. The layout of this paper is as follows. In next section we discuss our calculation of the SUSY virtual threshold effects. The final section contains discussions and conclusions of our numerical results.

II. VIRTUAL SUSY THRESHOLD EFFECTS

It is well-known that SUSY particles [gluinos, squarks] decrease or slow down the rate of fall of $\alpha_s(\mu)$ for large scale μ . “Large” means far above the threshold. At the one loop level the evolution equation for $\alpha_s(\mu)$ can be written as

$$\frac{d}{d\ln\mu}\alpha_s(\mu) = -\frac{b_3}{2\pi}\alpha_s^2(\mu) . \quad (1)$$

In the SM, b_3 is given by

$$b_3 = 11 - \frac{2}{3}n_f , \quad (2)$$

whereas in MSSM model one has

$$b_3 = 11 - \frac{2}{3}n_f - 2 - \frac{1}{3}\tilde{n}_f , \quad (3)$$

where n_f [$\tilde{n}_f$] is the number of quark [squark] flavors that are active. The contribution ‘−2’ is the gluino contribution [it is assumed that the gluino is active in this case]. In Ref. [4] the issue of the effect of high-mass thresholds due to gluinos, squarks and other new heavy quarks on the evolution of α_s was considered. The corrections to α_s were found to be appreciable, this in turn means a significant increase in the transverse momentum dependence of jet production at the Tevatron. However, as noted in Ref. [5], these authors [4] do not include the effect of $q\tilde{q}\tilde{g}$ Yukawa interactions and hence one cannot take their estimates for the superpartners of ordinary matter as final. In Ref. [5] the effect of Yukawa couplings was included. The results in [5] indicate that the CDF data cannot be explained by a mass threshold effect in the MSSM, as the calculated result is not only small but of the wrong sign. In a similar but more detailed analysis, the authors of Ref. [3], working in the context of MSSM, consider the virtual one-loop corrections to the parton-level subprocesses $q\bar{q} \rightarrow q\bar{q}$, $q\bar{q} \rightarrow q'\bar{q}'$, $qq' \rightarrow qq'$, $q\bar{q} \rightarrow gg$ and $qg \rightarrow qg$, which are expected to dominate the large p_T cross sections at the Tevatron energy.

The purpose of this talk is to give our results of incorporating the one-loop radiative corrections into the running of α_s , the dressing-up of the parton distribution functions, and finally convoluting the relevant subprocess cross sections with the SUSY dressed-up PDF’s. We note that one should take into account particles effects on the parton structure functions

at energies sufficiently far above the threshold and can ignore this effect in the threshold region. In the hadron colliders, like the Tevatron, what is measured is $p\bar{p}$ cross section, and not the individual subprocesses cross sections. So in order to determine the effect of subprocesses on the p_T cross section one must perform a convolution of the cross section of each subprocess with the corresponding PDF's. We find that in the process of convolution with the PDF's, the “dips and peaks” in the various subprocesses [3] are much reduced.

III. DISCUSSIONS AND CONCLUSIONS

For our numerical calculation, we implement various lower bounds on squarks and gluinos depending on parameters in the MSSM. For example, D0 group [6] searched for the events with large missing p_T with three or more jets, observed no such events above the level expected in the SM. This puts some limits on the squark and gluino masses assuming the short-lived gluinos:

$$m_{\tilde{g}} > 144 \text{ GeV} \quad \text{for} \quad m_{\tilde{q}} = \infty , \quad (4)$$

$$\text{or} \quad m_{\tilde{g}} = m_{\tilde{q}} > 212 \text{ GeV} . \quad (5)$$

CDF group is currently analyzing their data, with their preliminary data being similar to the D0 results with slight increase in sparticles' mass bound. In the subsequent numerical analysis, we choose three sets of $(m_{\tilde{g}}, m_{\tilde{q}})$, which we shall refer to as Case I, II and III, respectively,

$$(m_{\tilde{g}}, m_{\tilde{q}}) = (220 \text{ GeV}, 220 \text{ GeV}), \quad (150 \text{ GeV}, \infty), \quad (150 \text{ GeV}, 150 \text{ GeV}) . \quad (6)$$

The Case III is only of academic interest if the limit given in Eq. (5) is valid in reality.

Fig. 1 exhibits our results for $[(\frac{\text{MSSM}-\text{SM}}{\text{SM}}) \frac{d\sigma}{dp_{T_j}}]$ versus p_{T_j} for both Cases I and III [Case III refers the MSSM with the five squarks and gluinos being degenerate with a common mass

of 150 GeV, *i.e.* the last set in Eq. (6)]. A rapidity cut of 0.7 is used [$|\eta| < 0.7$]. A maximum difference of -1% from the SM is found for the both Cases. The question that naturally arises is, that why the “dips and peaks” [which are at the level of 5%–6% [3]] at the parton level, are reduced. There are two factors contributing to the reduction in the “dips and peaks” in the process of going from the parton to the process level. Part of this reduction comes from the process of convolution of the various subprocesses with PDF’s [as already remarked to in the previous section]. The other piece of this reduction in “dips and peaks” by going from subprocess to process level is due to t-channel “dilution” effect. This can be shown by simply not including the t-channel subprocesses’ contribution. When this is done [see Fig. 2] reduction in the “peaks and dips” is not so large.

In Fig. 2 we exhibit our results for $[(\frac{\text{MSSM}-\text{SM}}{\text{SM}})\frac{d\sigma}{dp_{T_j}}]$ versus p_{T_j} for both Cases I and III, but this time only including the subprocesses $q\bar{q} \longrightarrow q'\bar{q}'$ and $q\bar{q} \longrightarrow gg$. These two are the subprocesses with the prominent threshold structures from vacuum polarization interferences through s-channel exchange diagrams. A rapidity cut of 0.7 is used [$|\eta| < 0.7$]. The deviation from the SM for Case I varies between 2% to -4% , whereas as in Case II the variation is similar size to Case I at different energy scales related to the SUSY threshold effects.

In summary, in the MSSM the p_T distributions does not differ very much from those of the SM except for the possible threshold effects ($\sim 1\%$) through loop corrections. In actual experiments, the jet resolution will in general smear out any narrow resonance structures (which may be the case for the long-lived gluinos), leading to broad resonance structure, and therefore it looks impossible to detect SUSY particles through this kind of indirect virtual threshold effect. It has been reported [7] that the apparent discrepancy between CDF data and theory may be explained by the uncertainties resulting from the non-perturbative parton distribution, in particular in the gluon distribution. Our main conclusion seems to be on the right track in view of this global analysis of parton distribution function. For more details

on this talk, please look Ref. [8].

ACKNOWLEDGEMENTS

The work of CSK was supported in part by KOSEF, Project No. 951-0207-008-2, in part by CTP of SNU, in part by the BSRI Program, Project No. BSRI-97-2425, and in part by the COE Fellowship from Japanese Ministry of Education, Science and Culture.

REFERENCES

- [1] CDF Collaboration: F. Abe *et al.*, Phys. Rev. Lett. **77** (1996) 438.
- [2] D0 Collaboration: G. Blazey, talk given at Rencontre de Moriond, (March 1996).
- [3] J. Ellis and D. Ross, Phys. Lett. **B383** (1996) 187.
- [4] V. Barger *et al.*, Phys. Lett. **B383** (1996) 178.
- [5] P. Kraus and F. Wilczek, Phys. Lett. **B382** (1996) 262.
- [6] D0 Collaboration: S. Abachi *et al.*, Phys. Rev. Lett. **75** (1995) 618.
- [7] H.L. Lai *et al.*, hep-ph/9606399.
- [8] C.S. Kim and S. Alam, hep-ph/9610503.

FIGURES

Fig. 1 Deviation from the SM due to SUSY contribution to $\frac{d\sigma}{dp_{T_j}}$ versus p_{T_j} .

Fig. 2 Deviation from the SM due to SUSY contribution to $\frac{d\sigma}{dp_{T_j}}$ versus p_{T_j} , when only the subprocesses $q\bar{q} \rightarrow q'\bar{q}'$, and $q\bar{q} \rightarrow gg$ are included.




Fig. 1




Fig. 2